

13

# ON CONSTRUCTING SOME STRONGLY WELL-COVERED GRAPHS

by

Michael R. Pinter\*  
Department of Mathematics  
Belmont University  
Nashville, Tennessee, USA

AD-A261 848



DTIC  
ELECTE  
MAR 19 1993  
S B D

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

93-05542



16pgs

\* work partially supported by ONR Contracts #N00014-85-K-0488 and #N00014-91-J-1142

98 3 17 008

~~98 2 18 304~~

### Abstract

A graph is well-covered if every maximal independent set is a maximum independent set. If a well-covered graph  $G$  has the additional property that  $G-e$  is also well-covered for every line  $e$  in  $G$ , then we say the graph is strongly well-covered. We exhibit a construction which produces strongly well-covered graphs with arbitrarily large (even) independence number. The construction is in terms of a lexicographic graph product.

<b>Accession For</b>	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By <i>per letter</i>	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
<i>A-1</i>	

# ON CONSTRUCTING SOME STRONGLY WELL-COVERED GRAPHS

## INTRODUCTION

A set of points in a graph is independent if no two points in the graph are joined by a line. The maximum size possible for a set of independent points in a graph  $G$  is called the independence number of  $G$  and is denoted by  $\alpha(G)$ . A set of independent points which attains the maximum size is referred to as a maximum independent set. A set  $S$  of independent points in a graph is maximal (with respect to set inclusion) if the addition to  $S$  of any other point in the graph destroys the independence. In general, a maximal independent set in a graph is not necessarily maximum.

In a 1970 paper, Plummer [13] introduced the notion of considering graphs in which every maximal independent set is also maximum; he called a graph having this property a well-covered graph. The work on well-covered graphs that has appeared in the literature has focused on certain subclasses of well-covered graphs. Campbell [2] characterized all cubic well-covered graphs with connectivity at most two, and Campbell and Plummer [3] proved that there are only four 3-connected cubic planar well-covered graphs. Royle and Ellingham [16] have recently completed the picture for cubic well-covered graphs by determining all 3-connected cubic well-covered graphs.

For a well-covered graph with no isolated points, the independence number is at most one-half the size of the graph. Well-covered graphs whose independence number is exactly one-half the size of the graph are called very well-covered graphs. The subclass of very well-covered graphs was characterized by Staples [17] and includes all well-covered trees and all well-covered bipartite graphs. Independently, Ravindra [14] characterized bipartite well-covered graphs and Favaron [6] characterized the very well-covered graphs. Recently, Dean and Zito [4] characterized the very well-covered graphs as a subset of a more general (than well-covered) class of graphs.

A set  $S$  of points in a graph dominates a set  $V$  of points if every point in  $V-S$  is adjacent to at least one point of  $S$ . Finbow and Hartnell [7] and Finbow, Hartnell, and Nowakowski [8] studied well-covered graphs relative to the concept of dominating sets. Finbow, Hartnell, and Nowakowski have also obtained a characterization of well-covered graphs with girth at least five [9].

A well-covered graph is 1-well-covered if and only if the deletion of any point from the graph leaves a graph which is also well-covered. A well-covered graph is strongly well-covered if and only if the deletion of any line from the graph leaves a graph which is also well-covered. A well-covered graph is in the class  $W_2$  if and only if any two disjoint independent sets in the graph can be extended to disjoint maximum independent sets. Staples [18] showed that a well-covered graph is 1-well-covered if and only if it is in  $W_2$ . For the remainder of this paper, we use the  $W_2$  nomenclature instead of referring to 1-well-covered graphs.

The class of well-covered graphs contains all complete graphs and all complete bipartite graphs of the form  $K_{n,n}$ . The only cycles which are well-covered are  $C_3$ ,  $C_4$ ,  $C_5$ , and  $C_7$ . We note that all complete graphs (except  $K_1$ ) are also in  $W_2$ , but no complete bipartite graphs (except  $K_{1,1}$ ) are in  $W_2$ . The cycles  $C_3$  and  $C_5$  are the only cycles in  $W_2$ . Also note that the only complete graphs which are strongly well-covered are  $K_1$  and  $K_2$ , the only complete bipartite graphs which are strongly well-covered are  $K_{1,1}$  and  $K_{2,2}$ , and  $C_4$  is the only strongly well-covered cycle.

In [12], we show that a strongly well-covered graph with more than four points has minimum degree at least four and is 3-connected. Also, we show that all strongly well-covered graphs other than  $K_1$  and  $K_2$  have girth at most four, where the girth of a graph is the size of a smallest cycle in the graph and a graph with no cycles has infinite girth. In this paper we construct strongly well-covered graphs with triangles and strongly well-covered graphs with girth four.

## PRELIMINARY RESULTS

Unless otherwise stated, we assume all graphs are connected. Note that a disconnected graph is a  $W_2$  graph (strongly well-covered graph) if and only if each of its components is a  $W_2$  graph (strongly well-covered graph). For notation and terminology not defined here, see [1].

For a point  $v$  in a graph  $G$ , let  $N[v] = N(v) \cup \{v\}$ . Define  $G_v$  to be the graph induced by  $G - N[v]$ . In other words,  $G_v$  is the graph that remains after deleting  $v$  and all of its neighbors. In [12], the author shows that if  $G$  is a strongly well-covered graph and  $G$  is not complete, then for all points  $v$  in  $G$ , the graph  $G_v$  cannot contain a component which is a line. Campbell and Plummer [3] proved the following very useful necessary condition for a graph to be well-covered. We will use this later to verify a construction.

**Theorem 1.** If a graph  $G$  is well-covered and is not complete, then  $G_v$  is well-covered for all  $v$  in  $G$ . Moreover,  $\alpha(G_v) = \alpha(G) - 1$ .

Recall from earlier that if  $G$  is a  $W_2$  graph, then for all points  $v$  the graph  $G - v$  is well-covered (since a  $W_2$  graph is 1-well-covered). On the other hand, we show in [12] that strongly well-covered is a *sufficient* condition for  $G$  to have the property that for all points  $v$  the graph  $G - v$  is not well-covered. We state this here as Theorem 2. As a consequence,  $K_2$  is the only strongly well-covered graph which is also a  $W_2$  graph.

**Theorem 2.** If  $G$  ( $G \neq K_1$  or  $K_2$ ) is strongly well-covered, then for all points  $v$  in  $G$  the graph  $G - v$  is not well-covered.

Next, we state the characterization of the strongly well-covered graphs with independence number two, as given in [12]. This characterization will be quite helpful in building strongly well-covered graphs with independence number larger than two.

**Theorem 3.** Suppose  $G$  is well-covered with  $\alpha(G) = 2$ . Then  $G$  is strongly well-covered if and only if  $G$  is  $(|V(G)| - 2)$ -regular.

If  $G \neq K_2$  is well-covered and  $e = uv$  is a line in  $G$ , consider maximal independent sets in the graph  $G-e$ . Suppose  $J$  is a maximal independent set in  $G-e$  which does not contain at least one endpoint of  $e$  (that is,  $J \cap \{u, v\} \neq \{u, v\}$ ). Then it follows that  $J$  is a maximal independent set in  $G$ . Since  $G$  is well-covered, then  $|J| = \alpha(G)$ . Thus, every maximal independent set in  $G-e$  which does not contain at least one endpoint of  $e$  has size  $\alpha(G)$ . Consequently, to show that  $G-e$  is well-covered it suffices to show that every maximal independent set in the graph  $G-e$  which contains both endpoints of  $e$  has size  $\alpha(G)$ .

### A CONSTRUCTION

For our construction, we use a product of well-covered graphs. Suppose  $H$  is a graph with  $n$  points and  $\{G_i\}$ ,  $i = 1, \dots, n$ , is a family of disjoint graphs. Associate one member of  $\{G_i\}$  with each point of  $H$ . We assume  $V(H) = \{v_1, \dots, v_n\}$  and  $G_i$  is associated with  $v_i$ , for all  $i$ . We define the lexicographic product graph of  $H$  and  $\{G_i\}$ , denoted  $H \circ (G_1, \dots, G_n)$ , as follows:  $V(H \circ (G_1, \dots, G_n)) = V(G_1) \cup \dots \cup V(G_n)$  and  $E(H \circ (G_1, \dots, G_n)) = E(G_1) \cup \dots \cup E(G_n) \cup \{xy : x \in V(G_i), y \in V(G_j) \text{ and } v_i \sim v_j \text{ in } H\}$ .

If every member of the family  $\{G_i\}$  is the same graph  $G$ , then the lexicographic product consists of replacing each point of  $H$  with a copy of the graph  $G$  and joining the

copies as indicated above. In this special case, we denote the lexicographic product by  $H \circ G$ .

Topp and Volkmann [19] considered several different types of products of well-covered graphs. In particular for the lexicographic product of well-covered graphs, they proved a theorem which implies the following theorem.

**Theorem 4.** If  $H$  is well-covered and  $\{G_i\}$ ,  $i = 1, \dots, |V(H)|$ , is a family of well-covered graphs with  $\alpha(G_i) = \alpha(G_j)$  for all  $i$  and  $j$ , then  $H \circ (G_1, \dots, G_{|V(H)|})$  is well-covered. Moreover,  $\alpha(H \circ (G_1, \dots, G_{|V(H)|})) = \alpha(H)\alpha(G_1)$ .

In the next theorem, we give an additional condition on a well-covered graph  $H$  which is *sufficient* to obtain a strongly well-covered lexicographic product graph.

**Theorem 5.** Suppose  $H$  is a well-covered graph with the following additional property: if  $e = uv$  is a line in  $H$ , then  $H_{uv}$  is a well-covered graph and  $\alpha(H_{uv}) = \alpha(H) - 1$ , where  $H_{uv}$  is defined to be the graph  $H - (N[u] \cup N[v])$ .

Let  $|V(H)| = n$  and  $\{G_i\}$ ,  $i = 1, \dots, n$ , be a family of strongly well-covered graphs with  $\alpha(G_i) = 2$  for all  $i$  and each  $G_i$  is connected or  $2K_1$ . Let  $L = H \circ (G_1, \dots, G_n)$ . Then  $L$  is strongly well-covered, and  $\alpha(L) = 2\alpha(H)$ .

Proof. By Theorem 4, the lexicographic product graph  $L$  is well-covered and  $\alpha(L) = 2\alpha(H)$ . Let  $V(H) = \{u_1, u_2, \dots, u_n\}$ . Note the following about the structure of the lexicographic product graph:  $V(L)$  is the union of  $V(G_i)$ , for  $i = 1, \dots, n$ . If  $u_i \sim u_j$  in  $H$ , then  $x \sim y$  in  $L$  for all  $x \in V(G_i)$ , for all  $y \in V(G_j)$ . If  $u_i$  is not adjacent to  $u_j$  in  $H$  ( $i \neq j$ ), then  $x$  is not adjacent to  $y$  in  $L$  for all  $x \in V(G_i)$ , for all  $y \in V(G_j)$ . Also,  $a \sim b$  in  $G_i$  if and only if  $a \sim b$  in  $L$ , for all  $a$  and  $b$  in  $V(G_i)$ .

We proceed to show that  $L$  is *strongly* well-covered. Suppose  $e$  is a line in  $L$ . Then either  $e$  corresponds to a line in  $H$ , or  $e$  corresponds to a line in some  $G_j$ .

Case 1. Suppose  $e = xy$  corresponds to a line in  $G_j$ , for some  $j$ . Since  $G_j$  is strongly well-covered with  $\alpha(G_j) = 2$ , then  $\{x, y\}$  is a maximum independent set in the graph  $G_j - e$ . We consider the graph  $L - e$ .

To this end, consider the graph  $H_{u_j} = H - N[u_j]$  (a subgraph of  $H$ ). By Theorem 1, graph  $H_{u_j}$  is well-covered and  $\alpha(H_{u_j}) = \alpha(H) - 1$ . Let  $S_j$  be the subgraph of  $L$  corresponding to the components of  $H_{u_j}$ . Observe that  $S_j$  is a lexicographic product graph itself. Then since  $H_{u_j}$  is well-covered, by Theorem 4 the graph  $S_j$  is well-covered and  $\alpha(S_j) = 2\alpha(H_{u_j}) = 2(\alpha(H) - 1)$ .

Suppose  $J$  is a maximal independent set in  $L - e$  such that  $J \supseteq \{x, y\}$ . Since  $\{x, y\}$  is a maximum independent set in  $G_j - e$ , then  $J' = J - \{x, y\}$  must be contained in  $S_j$ . Since  $S_j$  is well-covered, each component of  $S_j$  is well-covered and it follows that  $|J'| = \alpha(S_j) = 2(\alpha(H) - 1)$ . Thus,  $|J| = 2\alpha(H)$ . So a maximal independent set in  $L - e$  which contains the endpoints of  $e$  has size  $2\alpha(H)$ . Thus, every maximal independent set in  $L - e$  has size  $2\alpha(H)$  and hence is a maximum independent set in  $L - e$ . Therefore,  $L - e$  is well-covered.

Case 2. Suppose  $e$  corresponds to the line  $u_i u_j$  in  $H$ . Say  $e = xy$ , where  $x \in V(G_i)$  and  $y \in V(G_j)$ .

By hypothesis,  $H_{u_i u_j}$  is well-covered and  $\alpha(H_{u_i u_j}) = \alpha(H) - 1$ . Suppose  $J \supseteq \{x, y\}$  is maximal independent in  $L - e$ . Let  $S_{ij}$  be the subgraph of  $L$  corresponding to  $H_{u_i u_j}$ . Observe that  $S_{ij}$  is a lexicographic product graph itself. Since  $H_{u_i u_j}$  is well-covered, then by Theorem 4 the graph  $S_{ij}$  is well-covered with  $\alpha(S_{ij}) = 2\alpha(H_{u_i u_j}) = 2(\alpha(H) - 1)$ . Let  $J' = J - \{x, y\}$ . Then  $J'$  is contained in  $S_{ij}$  and is maximal independent in  $S_{ij}$ . Thus,  $|J'| = 2(\alpha(H) - 1)$ , and so  $|J| = 2\alpha(H)$ . Hence, a maximal independent set in  $L - e$  which contains  $\{x, y\}$  necessarily has size  $2\alpha(H)$ . Since  $L$  is well-covered, then every maximal independent set in  $L - e$  has size  $2\alpha(H)$ . Thus,  $L - e$  is well-covered.

From Cases 1 and 2, we conclude that  $L - e$  is well-covered for all lines  $e$  in  $L$ . Therefore,  $L$  is strongly well-covered. []



Note in Theorem 5 that  $G_i$  is allowed to be disconnected. In this case,  $G_i$  must be  $2K_1$  since  $\alpha(G_i) = 2$ , the graphs  $K_1$  and  $K_2$  are the only complete strongly well-covered graphs, and from above, for every point  $v$  in  $G$ , the graph  $G_v$  cannot contain a component which is a line.

Although the condition in Theorem 5 is very restrictive, there are well-covered graphs which satisfy the condition and, hence, lead to the construction of infinite families of strongly well-covered graphs. We now give five such infinite families based on the five well-covered graphs shown in Figure 1.

**Corollary 6.** Suppose  $H$  is one of the five graphs in Figure 1 and  $\{G_i\}$ ,  $i = 1, \dots, |V(H)|$ , is a family of strongly well-covered graphs with  $\alpha(G_i) = 2$  and each  $G_i$  is connected or  $2K_1$ . Then  $H\circ(G_1, \dots, G_{|V(H)|})$  is strongly well-covered.

Proof. If  $H$  is one of the five graphs in Figure 1, it can be shown that  $H$  is well-covered, and for any line  $uv$  in  $H$ , the graph  $H_{uv} = H - (N[u] \cup N[v])$  is well-covered with  $\alpha(H_{uv}) = \alpha(H) - 1$ . By Theorem 5, it follows that  $H\circ(G_1, \dots, G_{|V(H)|})$  is strongly well-covered. []




Figure 1

We stated earlier that a strongly well-covered graph has girth at most four. From the following corollary, we are assured of the existence of strongly well-covered graphs with girth exactly four.

**Corollary 7.** If  $H$  is a triangle-free well-covered graph which satisfies the conditions in Theorem 5, then  $H \circ 2K_1$  is a girth 4 strongly well-covered graph.

Proof. If  $H$  is triangle-free, then  $H \circ 2K_1$  is also triangle-free. Clearly  $H \circ 2K_1$  has 4-cycles. The result then follows immediately from Theorem 5. []

For example, the graph in Figure 2 is  $C_5 \circ 2K_1$ . This graph was found by Royle [15] with the aid of a computer, and independently by the author.




Figure 2

## STRONGLY WELL-COVERED GRAPHS VIA $W_2$ GRAPHS OF GIRTH FOUR

From the graphs given in Figure 1, we can construct strongly well-covered graphs with  $\alpha \leq 8$ . In order to construct strongly well-covered graphs with arbitrarily large independence number, we turn to the family of  $W_2$  graphs of girth 4. First, we prove the following lemma about  $W_2$  graphs of girth 4, which will allow us to use Theorem 5 to construct families of strongly well-covered graphs.

**Lemma 8.** Suppose  $H$  is a  $W_2$  graph of girth 4 and  $e = uv$  is a line in  $H$ . Let  $H_{uv}$  be the graph  $H - (N[u] \cup N[v])$ . Then  $H_{uv}$  is well-covered and  $\alpha(H_{uv}) = \alpha(H) - 1$ .

**Proof.** Suppose  $e = uv$  is a line in  $H$ . Let  $U = N(u) - v$  and  $V = N(v) - u$ . Since  $H$  has no triangles, then  $U \cap V = \emptyset$ .

Suppose  $J$  is a maximal independent set in the graph  $H_{uv}$ . Clearly  $|J| < \alpha(H)$ . We wish to show that  $|J| = \alpha(H) - 1$ . We assume to the contrary that  $|J| < \alpha(H) - 1$ .

If  $J$  dominates  $V$ , then  $J \cup \{u\}$  is maximal independent in  $H$ . Since  $|J \cup \{u\}| < \alpha(H)$  and  $H$  is well-covered, we have a contradiction. Thus,  $J$  does not dominate  $V$ . Hence, there exists a point  $y$  such that  $y \in V$  and  $J$  does not dominate  $y$  (see Figure 3).




Figure 3

Note that  $N(y) - v$  is contained in  $V(H_{uv}) \cup U$ , since  $H$  has no triangles. Therefore,  $(J \cup \{u\}) \cap N(y) = \emptyset$ ,  $J \cup \{u\}$  is independent, and  $J \cup \{u\}$  dominates  $N(y)$ . It follows that  $J \cup \{u\}$  and  $\{y\}$  are disjoint independent sets in  $H$  which cannot be extended to disjoint maximum independent sets in  $H$ , and so  $H$  is not in  $W_2$ . This contradicts our hypothesis.

Thus,  $|J| = \alpha(H) - 1$ . Therefore, every maximal independent set in  $H_{uv}$  has size  $\alpha(H) - 1$ . It follows that  $H_{uv}$  is well-covered and  $\alpha(H_{uv}) = \alpha(H) - 1$ . []

In [10] and [11], the author presents constructions which yield  $W_2$  graphs of girth four with arbitrarily large independence number. Based on the  $W_2$  graphs obtained from these constructions, we show in the following theorem that we can construct infinite families of strongly well-covered graphs with arbitrarily large (even) independence number.

**Theorem 9.** Suppose  $H$  is a  $W_2$  graph of girth 4 with  $n$  points, and  $\{G_i\}$ ,  $i = 1, \dots, n$ , is a family of strongly well-covered graphs with  $\alpha(G_i) = 2$  and  $G_i$  is connected or  $2K_1$ , for all  $i$ . Then the lexicographic product graph  $H \circ (G_1, \dots, G_n)$  is a strongly well-covered graph, and  $\alpha(H \circ (G_1, \dots, G_n)) = 2\alpha(H)$ .

Proof. From Lemma 8, if  $e = uv$  is a line in  $H$ , then the graph  $H_{uv}$  is well-covered and  $\alpha(H_{uv}) = \alpha(H) - 1$ . Thus, the graph  $H$  satisfies the additional condition required of a well-covered graph in Theorem 5. It follows by Theorem 5 that  $H \circ (G_1, \dots, G_n)$  is strongly well-covered and  $\alpha(H \circ (G_1, \dots, G_n)) = 2\alpha(H)$ . []

Recall that a  $W_2$  graph  $H$  has the property that for all points  $v$  in  $H$ , the graph  $H-v$  is well-covered, and a strongly well-covered  $G$  has the property that for all points  $v$  in  $G$ , the graph  $G-v$  is *not* well-covered. Given this disparity between the two types of well-covered graphs, it is perhaps surprising that the lexicographic product of a  $W_2$  graph and a family of strongly well-covered graphs as produced in Theorem 9 will yield a strongly well-covered graph.

If  $H$  is a  $W_2$  graph of girth 4, then  $H \circ 2K_1$  is strongly well-covered by Theorem 9. Clearly,  $H \circ 2K_1$  has girth 4. Since there are infinitely many  $W_2$  graphs of girth 4, it follows that there are infinitely many *girth 4* strongly well-covered graphs.

The graphs given in Figure 4 are the strongly well-covered lexicographic product graphs  $H_1 \circ 2K_1$  and  $H_2 \circ 2K_1$ , where  $H_1$  and  $H_2$  are planar  $W_2$  graphs of girth 4 with eight points and eleven points, respectively (see [10] for a discussion of planar  $W_2$  graphs of

girth 4). Each of these graphs has points with degree four. Hence, the lower bound of four for the minimum degree in a strongly well-covered graph (mentioned above) is sharp.




Figure 4

A line in a graph  $G$  is a critical line if its removal increases the independence number. A line-critical graph is a graph with only critical lines. Staples proved in [17] that a triangle-free  $W_2$  graph is line-critical.

In searching for well-covered graphs  $H$  such that  $H_0(G_1, \dots, G_{|V(H)|})$  is strongly well-covered, for an appropriate family of graphs  $\{G_i\}$ , we discovered the following necessary condition on  $H$ .

**Theorem 10.** Suppose  $\{G_i\}$ ,  $i = 1, \dots, n$ , is a family of strongly well-covered graphs with  $\alpha(G_i) = 2$ , for all  $i$ . If  $H$  is a well-covered graph on  $n$  points and  $H_0(G_1, \dots, G_n)$  is strongly well-covered, then  $H$  is line-critical.

Proof. Assume to the contrary that  $e = uv$  is not a critical line in  $H$ . Thus,  $\alpha(H-e) = \alpha(H)$ . Let  $L = H_0(G_1, \dots, G_n)$ . Let  $e' = u_i v_j$  be a line in  $L$  corresponding to the line  $e$  in  $H$ , with  $u_i \in V(G_i)$ ,  $v_j \in V(G_j)$  ( $i \neq j$ ). Since  $\alpha(H-e) = \alpha(H)$ , then there exists a maximal independent set  $J$  in  $H-e$  which contains  $\{u, v\}$  such that  $|J| \leq \alpha(H)$ . So  $J - \{u, v\}$  dominates  $H_{uv}$  and is contained in  $V(H_{uv})$ . For  $x \in J - \{u, v\}$ , we have  $x \in V(G_m)$  for some  $m$ ,  $m \neq \{i, j\}$ . Since  $\alpha(G_m) = 2$ , there exists maximum independent set  $I_x \supset \{x\}$  in  $G_m$  with  $|I_x| = 2$ . Let  $I = \bigcup \{I_x : x \in J - \{u, v\}\}$ . So  $I$  is in  $V(L)$ . Since  $|J - \{u, v\}| \leq \alpha(H) - 2$  and  $|I_x| = 2$ , then  $|I| \leq 2(\alpha(H) - 2) = 2\alpha(H) - 4$ . But then  $I \cup \{u_i, v_j\}$  is maximal independent in  $L - e'$ , and  $|I \cup \{u_i, v_j\}| \leq 2\alpha(H) - 2 < 2\alpha(H)$ . Since  $\alpha(L) = 2\alpha(H)$  by Theorem 4 and  $L$  is assumed to be strongly well-covered, we have a contradiction. []

However, if  $H$  is line-critical, then  $H_0(G_1, \dots, G_{|V(H)|})$  is not necessarily strongly well-covered. In fact, being line-critical and in  $W_2$  are not *sufficient* conditions to ensure that  $H_0(G_1, \dots, G_{|V(H)|})$  is strongly well-covered. If  $H$  is the line-critical  $W_2$  graph in Figure 5, then  $H_0 K_1$  is not strongly well-covered. Note that the graph  $H_{uv} = H - (N[u] \cup N[v])$  is not well-covered.




Figure 5

## REFERENCES

1. J. A. Bondy and U. S. R. Murty, *Graph Theory With Applications* (American Elsevier, New York, 1976).
2. S. R. Campbell, Some results on planar well-covered graphs, *Ph.D. Dissertation*, Vanderbilt University, Nashville, TN, 1987.
3. S. R. Campbell and M. D. Plummer, On well-covered 3-polytopes, *Ars Combin.* 25-A (1988) 215-242.
4. N. Dean and J. Zito, Well-covered graphs and extendability, preprint, Bellcore, Morristown, NJ, and the The Johns Hopkins University, Baltimore, MD, 1990.
5. P. Erdős and T. Gallai, On the minimal number of vertices representing the edges of a graph, *Publ. Math. Inst. Hung. Acad. Sci.* 6 (1961) 181-203.
6. O. Favaron, Very well covered graphs, *Discrete Math.* 42 (1982) 177-187.
7. A. Finbow and B. Hartnell, On locating dominating sets and well-covered graphs, *Congr. Numer.* 65 (1988) 191-200.
8. A. Finbow, B. Hartnell, and R. Nowakowski, Well-dominated graphs: a collection of well-covered ones, *Ars Combin.* 25-A (1988) 5-10.
9. A. Finbow, B. Hartnell, and R. Nowakowski, A characterization of well-covered graphs of girth 5 or greater, to appear.
10. M. R. Pinter, A class of planar well-covered graphs with girth four, submitted, 1991.

11. M. R. Pinter, A class of well-covered graphs with girth four, submitted, 1991.
12. M. R. Pinter, Strongly well-covered graphs, submitted, 1991.
13. M. D. Plummer, Some covering concepts in graphs, *J. Combinatorial Theory* 8 (1970) 91-98.
14. G. Ravindra, Well-covered graphs, *J. Combin. Inform. System. Sci.* 2(1) (1977) 20-21.
15. G. F. Royle, Private communication to the author, 1991.
16. G. F. Royle and M. N. Ellingham, A characterization of well-covered cubic graphs, preprint, Vanderbilt University, Nashville, TN, 1991.
17. J. W. Staples, On some subclasses of well-covered graphs, *Ph.D. Dissertation*, Vanderbilt University, Nashville, TN, 1975.
18. J. W. Staples, On some subclasses of well-covered graphs, *J. Graph Theory* 3 (1979) 197-204.
19. J. Topp and L. Volkmann, On the well-coveredness of products of graphs, to appear.